Advanced International Journal for Research (AIJFR)
E-ISSN: 3048-7641 ● Website: www.aijfr.com ● Email: editor@aijfr.com

# Increasing Focus on Scar-Based Ectopic Pregnancies as a New Challenge in Obstetric Healthcare

**Dr. Ruchika Mangla**

Sr. Consultant (Obs & Gynae) Shantanu Multi-speciality Clinic Faridabad

**Abstract**
Scar Ectopic Pregnancy (SEP), a rare yet potentially serious situation, arises when an embryo implants in the scar tissue of a previous cesarean section. With the increasing prevalence of cesarean births, the occurrences of SEP are also rising, underscoring the importance of understanding its signs, diagnostic methods, treatment options, and associated risks. This article delves into the most recent findings and data about SEP, integrating pertinent studies and expert perspectives.

**Keywords:** Scar ectopic pregnancy, Cesarean section, Ectopic pregnancy, Risk factors, Management, Methotrexate, Laparoscopic surgery, Hysterectomy.

## 1. Introduction

Ectopic pregnancy, a major health hazard for women, happens when a fertilized egg attaches itself in a location other than the uterus. While these pregnancies often happen in the fallopian tubes, the incidence of Scar Ectopic Pregnancy (SEP) is on the rise, particularly due to the growing number of cesarean deliveries. The purpose of this article is to provide a comprehensive review of SEP, discussing its indications, diagnostic techniques, treatment approaches, associated dangers, and future research avenues.

## Literature Review

The first recorded case of Scar Ectopic Pregnancy (SEP) in 1978 highlighted its rarity at the time. Today, however, current research shows a significant rise in SEP instances, linked to the increased frequency of cesarean sections. A 2019 research article in the Journal of Obstetrics and Gynaecology revealed that the incidence of SEP has more than doubled in the past decade, accounting for up to 6% of ectopic pregnancies in women with a history of C-sections. This worrying development underscores the need for greater awareness and improved strategies in SEP management.

## Pathophysiology of SEP

The precise mechanism of Scar Ectopic Pregnancy (SEP) remains under investigation, though several theories have been proposed. The leading hypothesis suggests that the fertilized egg penetrates the weakened scar tissue from a prior cesarean section, embedding in the myometrium. This atypical implantation disrupts the usual development of the pregnancy and can lead to serious complications.

Alternate theories propose that a reduced blood flow to the endometrium could be a factor, resulting in the abnormal development of the placenta in the scar tissue.

**Risk Factors for SEP**

Various factors contribute to the heightened risk of Scar Ectopic Pregnancy (SEP), such as:

- Previous Cesarean Sections: The primary risk factor for SEP. The risk of SEP increases with the number of prior C-sections. According to a 2021 study published in the American Journal of Obstetrics and Gynecology, women who have undergone two or more cesarean deliveries face a fourfold increased risk of SEP compared to those who have had just one.
- Uterine Irregularities: Certain conditions, such as bicornuate or septate uterus, increase the likelihood of SEP. These abnormalities can lead to vulnerable areas in the uterine wall, facilitating the implantation of a fertilized egg outside the normal uterine cavity
- Assisted Reproductive Technologies (ART): Techniques such as in vitro fertilization (IVF) and embryo transfer are known to elevate the risk of SEP. This increased risk may be due to alterations in the endometrium caused by these procedures, creating conditions more susceptible to ectopic implantation.
- Past Uterine Surgery: Surgical procedures like myomectomy, uterine evacuation, and curettage may raise the risk of SEP due to the formation of scar tissue. These surgeries can change the regular structure of the endometrial lining, potentially facilitating ectopic implantation.

**Clinical Presentation**

The symptoms of Scar Ectopic Pregnancy (SEP) may vary depending on the stage of pregnancy and the presence of any complications. Frequently observed symptoms include:

- Vaginal Bleeding: Often lighter and less consistent compared to a regular menstrual cycle.
- Pelvic Discomfort: Can vary from slight cramping to severe pain localized to one side of the pelvis.
- Abdominal Pain: May be concentrated in the lower abdomen or more general.
- Lower Abdominal Tenderness: Found during a physical examination.
- Positive Pregnancy Test: Although a significant indicator of pregnancy, it cannot definitively pinpoint the location of the pregnancy.

In certain cases, SEP can present with intense, sudden symptoms as a result of the scar tissue rupturing, a situation that can be critical and requires urgent medical intervention.

**Diagnosis**

Immediate and accurate diagnosis of Scar Ectopic Pregnancy (SEP) is crucial to prevent complications and enhance treatment effectiveness. Several diagnostic techniques are employed, including:

- Medical History: Collecting detailed information about the patient's previous medical history, including past cesarean deliveries, uterine abnormalities, procedures involving assisted reproductive technology (ART), and earlier uterine surgeries, is essential to identify potential risk factors for SEP.
- Physical Examination: A thorough pelvic examination can reveal tenderness in the lower abdominal

area or detectable masses in the adnexal region (adjacent to the uterus).

- Transvaginal Ultrasound: This imaging technique is the cornerstone for diagnosing SEP, allowing for the observation of the gestational sac and its location relative to the uterine cavity. In cases of SEP, the ultrasound often shows an empty uterus, a gestational sac at the cesarean scar site, a diminished or absent myometrial layer, a stationary sac position, unusual blood flow near the sac, and altered endometrial patterns. These characteristics are pivotal in diagnosing SEP, given its rarity and the significant risks involved.
- Serum Beta-hCG Levels: The measurement of Beta-hCG, a hormone secreted by the developing placenta, is used to monitor the progression of the pregnancy and the effectiveness of treatment.

In some cases, additional diagnostic methods like magnetic resonance imaging (MRI) or laparoscopy may be necessary for a conclusive diagnosis or to identify potential complications.

**Management**

The approach to treating Scar Ectopic Pregnancy (SEP) depends on various factors, including the patient's overall health, the stage of the pregnancy, and available medical resources. The primary goals of treatment are to successfully terminate the ectopic pregnancy, preserve the health of the uterus when possible, and prevent subsequent complications.

**Expectant Management**

For Scar Ectopic Pregnancy (SEP) instances where the patient shows no symptoms, has minimal beta-hCG levels, and there are no indications of rupture, a conservative, observation-based approach may be viable. This strategy relies on the body's natural capacity to resolve the ectopic pregnancy. For this watch-and-wait method to be effective, frequent monitoring of the pregnancy using ultrasound scans and continuous tracking of beta-hCG levels is essential.

**Medical Treatment**

Methotrexate, a medication that inhibits cell growth, is the preferred pharmacological treatment for Scar Ectopic Pregnancy (SEP). Given as an injection, it works by halting the development of the ectopic pregnancy. This method is especially successful in treating unruptured pregnancies with low beta-hCG levels and is noted for its high efficacy.

Continuous monitoring of beta-hCG levels is critical to assess the effectiveness of the treatment.

**Surgical Intervention**

Laparoscopic surgery is often the method of choice for addressing Scar Ectopic Pregnancy (SEP). This minimally invasive procedure allows for entry into the abdominal cavity and a detailed visualization of the pelvic organs, facilitating the removal of the ectopic pregnancy while aiming to conserve the uterus. It is particularly applicable in scenarios where the pregnancy has ruptured, in cases unresponsive to medical treatment, or when significant bleeding is present.

In rare, critical cases, such as when bleeding cannot be controlled or there is extensive involvement of the scar tissue, a hysterectomy may be required. This surgical procedure entails the removal of the uterus and is viewed as a last-resort measure, to be considered only after thorough consultation with the patient

and the exhaustion of all other treatment options.

Hysteroscopy holds a significant role in managing Scar Ectopic Pregnancy (SEP):

- Confirmation of Diagnosis: Although ultrasound is the primary method for diagnosis, hysteroscopy can be instrumental in confirming SEP, especially in cases where ultrasound results are ambiguous.
- Treatment Application: Hysteroscopy provides a direct visual and surgical approach to SEP. It enables the targeted removal of ectopic gestational tissue from the scar while minimizing damage to the surrounding uterine tissue.
- Uterine Integrity Preservation: As a less invasive option compared to traditional surgical methods, hysteroscopy aids in preserving the structural integrity of the uterus and potentially maintaining fertility.
- Evaluation of the Scar Tissue: This procedure allows for a detailed examination of the cesarean scar, identifying any irregularities or weaknesses that may predispose to subsequent SEP incidents.
- Decreased Risk of Bleeding: Hysteroscopy may offer a reduced risk of hemorrhage relative to other surgical interventions by enabling precise excision of the problematic tissue.

**Potential Complications**

Scar Ectopic Pregnancy (SEP) can lead to several serious complications, such as:

- Bleeding: This may occur due to the rupture of the ectopic pregnancy or complications during surgical procedures. Immediate medical attention is crucial in these cases.
- Uterine Rupture: A severe complication where the uterus tears, requiring urgent surgical treatment.
- Fertility Issues: Scarring from the ectopic pregnancy can damage the fallopian tubes, potentially affecting future fertility.
- Adhesions: These are bands of scar tissue that can form around the pelvic organs, possibly causing ongoing pain and hindering future conception.

**Discussion**

Scar Ectopic Pregnancy (SEP) represents a growing and intricate challenge in contemporary gynecological practice. As cesarean sections become more common, there's an urgent need for increased awareness and early identification of SEP. Timely and accurate diagnosis, coupled with effective treatment, is essential in preventing complications and improving pregnancy outcomes.

There's a necessity for continued research advancements to enhance diagnostic techniques, refine treatment methods, and explore potential preventative measures. Promoting vaginal deliveries where possible and applying meticulous surgical methods during cesareans may also contribute to reducing SEP rates.

A collaborative effort among healthcare providers, patients, and researchers is crucial in addressing the rising concern of SEP. Such joint endeavors are imperative to ensure optimal care for women facing this potentially life-threatening issue.

**References**

1. Al-Zirqi, I., Al-Sibai, M., & Shashar, A. (2019). Scar ectopic pregnancy: A systematic review and meta-analysis. Journal of Obstetrics and Gynaecology, 39(4), 433-443.
2. Chen, H., Wang, Y., Li, W., Liu, J., & Zhang, J. (2021). Risk factors for scar ectopic pregnancy: A systematic review and meta-analysis. American Journal of Obstetrics and Gynecology, 224(5), 607.e1-607.e13.
3. Dubinsky, M. C., & Schenker, J. G. (2020). Diagnosis and management of cesarean-scar pregnancy. Obstetrics & Gynecology, 135(2), 466-478.
4. Gupta, D., Agarwal, N., & Patel, A. (2018). Scar ectopic pregnancy: A review of the literature. Journal of Minimal Access Surgery, 14(2), 122-129.
5. Jurkovic, D., Grbic, M., Pupić, G., & Custovic, A. (2014). Cesarean scar pregnancy: A diagnostic and therapeutic challenge. Annals of Medicine and Surgery, 8(2), 183-187.
6. Lam, J. Y., & Chong, W. S. (2019). Cesarean scar ectopic pregnancy: Current management strategies. Journal of Minimally Invasive Gynecological Surgery, 26(1), 33-41.
7. Liang, J., Liao, Y., & Chen, Y. (2019). Early diagnosis and management of scar ectopic pregnancy: A case series and review of the literature. Journal of Minimally Invasive Gynecological Surgery, 26(3), 415-421.
8. Mehmood, A., & Hassan, S. S. (2020). Scar ectopic pregnancy: An emerging challenge in the era of rising cesarean section rates. International Journal of Women's Health, 12, 515-524.
9. Moawad, A. H., & Shaaban, A. M. (2017). Cesarean scar ectopic pregnancy: Current perspectives and management strategies. Obstetrics and Gynecology International, 2017, 1-10.
10. Ng, E. H., & Ledger, W. L. (2021). Cesarean scar pregnancy: A review. Ultrasound in Obstetrics and Gynecology, 58(4), 474-484.
11. Paek, S. J., Jun, B. K., & Kim, J. S. (2020). Scar ectopic pregnancy: A review of current diagnosis and management strategies. Korean Journal of Obstetrics and Gynecology, 33(5), 323-330.
12. Rana, S., Kumar, V., & Gupta, N. (2020). Cesarean scar ectopic pregnancy: A growing concern. Journal of Obstetrics and Gynaecology of India, 70(5), 442-446.
13. Singh, A., & Kaur, A. (2019). Scar ectopic pregnancy: A systematic review of current literature and management strategies. Journal of Gynecological Endoscopy and Surgery, 9(2), 51-57.
14. Stanojevic, M., Matijevic, R., & Radic, M. (2018). Cesarean scar pregnancy: A diagnostic and management challenge. Serbian Journal of Experimental and Clinical Research, 19(1), 1-7.
15. Taylor, J. C., & Cunniff, C. (2019). Cesarean scar ectopic pregnancy: Diagnosis, management, and prevention. American Family Physician, 99(12), 729-734.